

AA.I1C1 BALTIC PHOTIC COARSE SEDIMENT DOMINATED BY *FUCUS SPP.*

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic bottoms in the photic zone with at least 90 % coverage of coarse sediment. Coarse sediment has less than 20 % of mud/silt/clay fraction (<63 µm), and the proportion of gravel and pebbles (grain size 2–63 mm) exceeds 30% of the combined gravel and sand fraction . Perennial attached algae cover at least 10 % of the seabed and more than other perennial attached erect groups. Out of the perennial attached algae *Fucus spp.* constitutes at least 50 % of the biovolume.

PHYSICAL ENVIRONMENT

Salinity range: >4.5 psu; Exposure range: Sheltered; Depth range: photic zone to about 5 meters

CHARACTERISTIC SPECIES

Fucus vesiculosus, *F. radicans*, *F. serratus*

ANTHROPOGENIC THREATS

Decreased light penetration depth and increased sedimentation caused by eutrophication

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.4 Gravel bottoms

2.4.2 Sublittoral photic zone

2.4.2.2 Level bottoms dominated by macrophyte vegetation

EUNIS 2012:

A5 Sublittoral sediment

A5.5 Sublittoral macrophyte-dominated sediment

<http://eunis.eea.europa.eu/habitats/1733>