

AA.I2 BALTIC PHOTIC COARSE SEDIMENT CHARACTERIZED BY SPARSE MACROSCOPIC EPIBENTHIC BIOTIC STRUCTURES

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic bottoms in the photic zone with at least 90 % coverage of coarse sediment. Coarse sediment has less than 20 % of mud/silt/clay fraction (<63 µm), and the proportion of gravel and pebbles (grain size 2–63 mm) exceeds 30 % of the combined gravel and sand fraction . Coverage of macroscopic vegetation or sessile macroscopic epifauna is 0><10 %.

PHYSICAL ENVIRONMENT

Salinity range: all; Exposure range: moderate; Depth range: photic zone

CHARACTERISTIC SPECIES

Mytilus spp., *Macoma baltica*, *Bathyporeia pilosa*, *Hediste diversicolor*

MAPPING ADVISE (HABITAT DELINEATION, IDENTIFICATION, SIMILAR TYPES)

Photic zone areas with coarse sediment such as gravel. Sediment must contain less than 20 % of silt, clay or mud, and at least 30 % of grain size 2–63 mm. Coverage of macroscopic vegetation or sessile macroscopic epifauna is 0><10 %.

GEOGRAPHIC RANGE

Whole Baltic Sea

ANTHROPOGENIC THREATS

Silting caused by eutrophication, dredging spoil deposition etc.

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.4 Gravel bottoms

2.4.2 Sublittoral photic zone

2.4.2.1 Level bottoms with little or no macrophyte vegetation

EUNIS 2012:

A5 Sublittoral sediment

A5.1 Sublittoral coarse sediment

A5.11 Infralittoral coarse sediment in low or reduced salinity

A5.111 : Baltic level gravel bottoms of the infralittoral photic zone with little or no macrophyte vegetation

<http://eunis.eea.europa.eu/habitats/2576>