

# AA.H1K1 BALTIC PHOTIC MUDDY SEDIMENT DOMINATED BY TUBE-BUILDING POLYCHAETES

## AUTHOR

HELCOM RED LIST Biotope Expert Team

## TEXTUAL DESCRIPTION

Baltic photic zone bottoms with at least 90 % coverage of muddy sediment. The sediment must contain at least 20 % of mud, silt or clay (grain size less than 63 µm). Epibenthic polychaetes cover at least 10% of the seabed and more than other perennial attached erect groups. Out of the epibenthic polychaetes tube-building polychaetes constitute at least 50 % of the biomass.

## PHYSICAL ENVIRONMENT

Substrate is muddy sediment. Appears in low to moderate energy exposure classes.

## CHARACTERISTIC SPECIES

Several species from the taxa Maldanidae and Terebellida.

## QUALITY DESCRIPTORS

Diversity, abundance and biomass of fauna.

## GEOGRAPHIC RANGE

Western Baltic Sea, including the Sound and Kattegat.

## ANTHROPOGENIC THREATS

Eutrophication, contaminants

## CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

### HELCOM 1998:

2.7 Muddy bottoms

2.7.2 Sublittoral photic zone

2.7.2.1 With little or no macrophyte vegetation

**EUNIS 2012:**

A5 Sublittoral sediment

A5.3 Sublittoral mud

A5.31 Sublittoral mud in low or reduced salinity

A5.311 Baltic brackish water sublittoral muddy biocenoses influenced by varying salinity

<http://eunis.eea.europa.eu/habitats/2585>