

AA.H3L8 BALTIC PHOTIC MUDDY SEDIMENT DOMINATED BY ABRA SPP.

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic photic zone bottoms with at least 90 % coverage of muddy sediment. The sediment must contain at least 20 % of mud, silt or clay (grain size less than 63 µm). Biomass of infaunal bivalves dominates and is highest in the group that includes infaunal bivalves/polychaetes/crustaceans/echinoderms/insect larvae. Out of the infaunal bivalves *Abra* spp. constitutes at least 50 % of the biomass.

PHYSICAL ENVIRONMENT

Substrate is muddy sediment. Appears in low to moderate energy exposure classes.

CHARACTERISTIC SPECIES

Abra spp.

QUALITY DESCRIPTORS

Diversity, abundance and biomass of fauna.

ANTHROPOGENIC THREATS

Eutrophication

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.7 Muddy bottoms

2.7.2 Sublittoral photic zone

2.7.2.1 With little or no macrophyte vegetation

EUNIS 2012:

A5 Sublittoral sediment

A5.3 Sublittoral mud

A5.31 Sublittoral mud in low or reduced salinity

A5.311 Baltic brackish water sublittoral muddy biocenoses influenced by varying salinity

<http://eunis.eea.europa.eu/habitats/2585>