

# AA.J3N BALTIC PHOTIC SAND CHARACTERIZED BY INFAUNAL CRUSTACEA

## AUTHOR

HELCOM RED LIST Biotope Expert Team

## TEXTUAL DESCRIPTION

Baltic photic bottoms with at least 90 % coverage of sand. Sand has less than 20 % of mud/silt/clay fraction (<63 µm), and the proportion of sand (grain size 0.063–2 mm) exceeds 70% of the combined gravel and sand fraction. No macrovegetation or epibenthic macrofauna. Biomass of infaunal crustacea dominates in the group that includes infaunal bivalves/polychaetes/crustaceans/insect larvae.

## PHYSICAL ENVIRONMENT

Substrate is sand. Appears in high energy exposure areas.

## CHARACTERISTIC SPECIES

*Monoporeia affinis*

## QUALITY DESCRIPTORS

Diversity, abundance and biomass of fauna.

## GEOGRAPHIC RANGE

Whole Baltic Sea

## CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

### **HELCOM 1998:**

2.5 Sandy bottoms

2.5.2 Sublittoral photic zone

2.5.2.1 Level bottoms with little or no macrophyte vegetation

### **EUNIS 2012:**

A5 Sublittoral sediment

A5.2 Sublittoral sand

A5.21 Sublittoral sand in low or reduced salinity

A5.211 Baltic level sandy bottoms of the infralittoral photic zone with little or no macrophyte vegetation

<http://eunis.eea.europa.eu/habitats/2580>