

AA.J3N3 BALTIC PHOTIC SAND DOMINATED BY SAND DIGGER SHRIMP (*BATHYPOREIA PILOSA*)

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic photic zone bottoms with at least 90 % coverage of sand. Sand has less than 20 % of mud/silt/clay fraction (<63 µm), and the proportion of sand (grain size 0.063–2 mm) exceeds 70% of the combined gravel and sand fraction. Biomass of infaunal crustaceans dominates and is highest in the group that includes infaunal bivalves/polychaetes/crustaceans/echinoderms/insect larvae. Out of the infaunal crustacean, *Bathyporeia pilosa* constitutes at least 50 % of the biomass

PHYSICAL ENVIRONMENT

Substrate is sand. Depth is typically from 1 to 10 meters, Appears in areas of moderate to high wave exposure. Salinity >4 psu.

CHARACTERISTIC SPECIES

Bathyporeia pilosa, *Oligochaeta*, *Haustorius arenarius*, *Cyathura carinata*, *Hediste diversicolor*

QUALITY DESCRIPTORS

Diversity, abundance and biomass of fauna.

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.5 Sandy bottoms

2.5.2 Sublittoral photic zone

2.5.2.1 Level bottoms with little or no macrophyte
vegetation

EUNIS 2012:

A5 Sublittoral sediment

A5.2 Sublittoral sand

A5.21 Sublittoral sand in low or reduced salinity

A5.211 Baltic level sandy bottoms of the infralittoral photic zone with little or no macrophyte vegetation

<http://eunis.eea.europa.eu/habitats/2580>