

AA.M1B3 BALTIC PHOTIC MIXED SUBSTRATE DOMINATED BY WATERMILFOIL (*MYRIOPHYLLUM SPICATUM* AND/OR *MYRIOPHYLLUM SIBIRICUM*)

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic bottoms in the photic zone with more than 10 %, but less than 90 % coverage of both hard and soft substrata. Submerged rooted plants cover least 10 % of the seabed and more than other perennial attached erect groups. Out of the submerged rooted plants, watermilfoil constitutes at least 50 % of the biovolume.

PHYSICAL ENVIRONMENT

Salinity range: <6 psu; Exposure range: low; Depth range: photic zone from about 0.2 to 2 meters.

CHARACTERISTIC SPECIES

Myriophyllum spicatum, *Myriophyllum sibiricum*

GEOGRAPHIC RANGE

Whole Baltic Sea

ANTHROPOGENIC THREATS

Eutrophication

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.8 Mixed sediment bottoms

2.8.2 Sublittoral photic zone

2.8.2.2 Dominated by macrophyte vegetation

EUNIS 2012:

A5 Sublittoral sediment

A5.5 Sublittoral macrophyte-dominated sediment

A5.54 Angiosperm communities in reduced salinity

<http://eunis.eea.europa.eu/habitats/1591>