

AB.E1F1 BALTIC APHOTIC SHELL GRAVEL DOMINATED BY VASE TUNICATE (*CIONA INTESTINALIS*)

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic aphotic bottoms at least 90 % coverage of shell gravel. Epibenthic chordates cover at least 10% of the seabed and more than other perennial attached erect groups. Out of the epibenthic chordates, vase tunicate constitutes at least 50 % of the biomass.

PHYSICAL ENVIRONMENT

Substrate is shell gravel.

CHARACTERISTIC SPECIES

Ciona intestinalis

GEOGRAPHIC RANGE

Known from German waters in the Baltic Sea

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.6 Shell gravel bottoms

2.6.1 Aphotic zone

HELCOM 2007:

Shell gravel bottoms

- habitat under threat and/or in decline in all areas of occurrence

The Southern Baltic Proper, The Gulf of Gdansk, Bay of Mecklenburg, Kiel Bay, Little Belt, Great Belt, The Sound, Kattegat

EUNIS 2012:

A5 Sublittoral sediment

A5.1 Sublittoral coarse sediment

A5.11 Infralittoral coarse sediment in low or reduced salinity

A5.115 Baltic shell gravel bottoms of the aphotic zone