

AB.E4 BALTIC APHOTIC SHELL GRAVEL CHARACTERIZED BY NO MACROSCOPIC BIOTIC STRUCTURES

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic aphotic bottoms with at least 90 % coverage of shell gravel. No macro- epi- or infauna.

PHYSICAL ENVIRONMENT

Substrate is shell gravel. Appears mostly in high energy exposure areas.

MAPPING ADVISE (HABITAT DELINEATION, IDENTIFICATION, SIMILAR TYPES)

No macro- epi- or infauna

GEOGRAPHIC RANGE

Southern part of Baltic Sea, the Sound, Kattegat

ANTHROPOGENIC THREATS

Increase in atmospheric CO₂ (Ocean acidification)

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.6 Shell gravel bottoms

2.6.1 Aphotic zone

HELCOM 2007:

Shell gravel bottoms

- habitat under threat and/or in decline in all areas of occurrence: The Southern Baltic Proper, The Gulf of Gdansk, Bay of Mecklenburg, Kiel Bay, Little Belt, Great Belt, The Sound, Kattegat

EUNIS 2012:

A5 Sublittoral sediment

A5.1 Sublittoral coarse sediment

A5.11 Infralittoral coarse sediment in low or reduced salinity

A5.115 Baltic shell gravel bottoms of the aphotic zone