

AB.J3L4 BALTIC APHOTIC SAND DOMINATED BY SAND GAPER (*MYA ARENARIA*)

AUTHOR

HELCOM RED LIST Biotope Expert Team

TEXTUAL DESCRIPTION

Baltic aphotic zone bottoms with at least 90 % coverage of sand. Sand has less than 20 % of mud/silt/clay fraction (<63 µm), and the proportion of sand (grain size 0.063–2 mm) exceeds 70% of the combined gravel and sand fraction. Biomass of infaunal bivalves dominates and is highest in the group that includes infaunal bivalves/polychaetes/crustaceans/echinoderms/insect larvae. Out of the infaunal bivalves, *Mya arenaria* constitutes at least 50 % of the biomass of the infaunal bivalves.

CHARACTERISTIC SPECIES

Mya arenaria

QUALITY DESCRIPTORS

Diversity, abundance and biomass of fauna

GEOGRAPHIC RANGE

Known from German waters in the Baltic Sea

CORRESPONDENCE WITH OTHER CLASSIFICATION SYSTEMS

HELCOM 1998:

2.5 Sandy bottoms

2.5.1 Aphotic zone

EUNIS 2012:

A5 Sublittoral sediment

A5.2 Sublittoral sand

A5.27 Deep circalittoral sand

A5.273 Baltic sandy bottoms of the aphotic zone

<http://eunis.eea.europa.eu/habitats/2620>